

RESEARCH REPORT

PERLITE AND
POLYSTYRENE
GROWING MEDIA


PERLITE PLANT GUIDE

The Perlite Institute Inc.
1924 North Second Street
Harrisburg, PA 17102 USA
tel. 717-238-9723 fax. 717-238-9985
Email: info@perlite.org

COMPARATIVE GROWTH STUDIES PERLITE VS. POLYSTYRENE MEDIA

by
O. A. Matkin
Soil and Plant Laboratory, Inc.
Santa Ana, California

Previous measurements of physical properties of growing media comparing perlite with polystyrene beads have indicated higher density and moisture retention when perlite is used in the growing media. Since high moisture holding capacity combined with adequate aeration is an important asset in container production of plants, trials were carried out comparing ryegrass growth in mixes of watering schedules. Composition of the two amendment mixes is shown in Table 1.



Photograph comparing ryegrass grown in perlite mix on left and polystyrene bead mix on right. Perlite mix C and polystyrene bead mix C were watered every three days beginning with the second week after seeding.

Table 1 Composition of Growing Media

Perlite Media	Polystyrene Media
0.5 cu. Yds. Perlite 0.5 cu. Yds. Sphagnum peat moss 5 lbs. Calcium carbonate lime 3 lbs. Dolomite lime	0.5 cu. Yds. Perlite 0.5 cu. Yds. Sphagnum peat moss 5 lbs. Calcium carbonate lime 3 lbs. Dolomite lime

Watering Schedule Varied

Containers used in growth trials were 6 inch plastic pots, 20 for each mix, 1400 milliliters of mix per pot. All pots were seeded to perennial ryegrass at the same time and maintained moist for 1 week at which time uniform germination had occurred. Beginning with the second week, the watering schedule was varied according to the schedule shown in Table 2.

The water that was supplied contained a balanced soluble nutrient and was supplied to each pot in the amount of 500 milliliters per pot at each scheduled irrigation. The excess liquid from each pot was collected and returned to the pot and drainage water was again collected, volume measured and recorded before discarding.

Perlite A Polystyrene A	Every Day
Perlite B Polystyrene B	Every 2 Days
Perlite C Polystyrene C	Every 3 Days
Perlite D Polystyrene D	Every 4 Days


"...SAMPLES PLANTED IN POLYSTYRENE BEAD MIX AND WATERED ONCE EVERY 4 DAYS FAILED TO PRODUCE ADDITIONAL GROWTH AFTER 12 WEEKS."

Clippings Weighed

The use of ryegrass as the test crop permitted successive harvests at which time clippings were dried and weighed. Harvests were made at 4, 8, 12, 16 and 20 weeks

The maximum volume of water retained after drainage was determined to be 583 milliliters for the peat/perlite mix and 383 milliliters for the peat/polystyrene bead mix (1400 milliliters volume of mix).

During the course of the growth trials, some unusually high temperatures occurred in the greenhouses which placed severe stress on crops under the limited irrigation regime. The five ryegrass samples planted in a polystyrene bead mix and watered once every 4 days failed to produce additional growth after 12 weeks.



Accumulative yield of clippings of ryegrass grown in perlite media and polystyrene media at four different irrigation schedules.

Results

Growth, as measured by quantity of clipping yield (see curve), was best in most frequently water samples. Visual inspection confirmed parallel root development. The order of growth rating from highest to lowest was: Perlite A, Perlite B, Polystyrene A, Perlite C, Polystyrene B, Perlite D, Polystyrene C,

